

Whispering gallery mode enhanced optical force with resonant tunneling excitation in the Kretschmann geometry

J. J. Xiao,^{1,*} Jack Ng,^{1,†} Z. F. Lin,² and C. T. Chan¹

¹Department of Physics and William Mong Institute of NanoScience and Technology,

The Hong Kong University of Science and Technology, Clear Water Bay, Hong Kong, China

²Surface Physics Laboratory and Department of Physics, Fudan University, Shanghai 200433, China

(Dated: January 5, 2009)

The boundary element method is applied to investigate the optical forces when whispering gallery modes (WGMs) are excited by a total internally reflected wave. Such evanescent wave is particularly effective in exciting the high- Q WGM, while the low angular or high radial order modes are suppressed relatively. This results in a large contrast between the forces on and off resonance, and thus allows for high size-selectivity. We fully incorporate the prism-particle interaction and found that the optical force behaves differently at different separations. Optimal separation is found which corresponds to a compromise between intensity and Q factor.

Microcavities with whispering gallery mode (WGM) are important components in nanophotonics, nonlinear optics, quantum optics, and many other areas.¹ As it is very difficult to accurately control the resonance frequency or the size of a micro-cavity during its fabrication, a convenient way to pick up the particles with the desired resonance frequencies or size is highly desirable. Recently, it was proposed in Ref. 2 that the optical force acting on a microsphere under an evanescent wave illumination^{3,4,5,6,7} is highly size-selective: at resonance the microsphere experiences a sizable force, whereas at off resonance the force is negligible.⁸ Utilizing this property and the high quality factor ($Q > 10^7$) of WGM,^{1,9} the size-selective force would allow for accurate size-sorting of microsphere cavities, with an accuracy of $\sim 1/Q$. However, the generation of evanescent wave requires a substrate. Ref. 2 considers only the regime where the microsphere is away from the near field of the substrate, and the interaction of the microsphere and substrate is only treated in a phenomenological manner.

In this article, using the boundary element method (BEM)¹⁰ and the Maxwell stress tensor,¹¹ we intend to go beyond Ref. 2 by fully incorporating the influence of the substrate. We confine ourselves to 2D calculations with a transverse electric (TE) polarization, since 3D calculation is difficult due to the need of extremely fine meshes in the BEM calculation. We expect that our result should be qualitatively relevant for the 3D system. We consider the Kretschmann geometry as depicted in Fig. 1(f): a microcylinder (MC) of radius a sits on top of a prism, both have a refractive index of $n_p = 1.5$. The combined system (in air) is then illuminated by an incident plane wave propagating at $\theta = 45^\circ$. As the critical angle of the prism is $\theta_c = 41.8^\circ$, the incident wave is total internally reflected at the prism surface, and an evanescent wave is generated there. The transfer of photon momentum to the MC is allowed by the scattering or tunneling of the evanescent wave. We shall see that when the particle is not in the near field of the prism surface, the treatment of Ref. 2 gives a correct description of the situation. For the near field case, we observe a series of interesting phenomena, some are caused by the interplay

between the increasing intensity and the decreasing Q as the particle approaches the surface, and some are caused by the interaction between the particle and the substrate.

The research on WGM-induced forces^{12,13,14,15,16,17,18} can be dated back to the early eighties when Ashkin reported the first observation of the force in air.¹⁹ Later, the WGM-induced forces were also observed by Fontes *et al* in water.²⁰ Nevertheless, only a modest enhancement in the optical force has been achieved owing to the use of propagating waves in these works. Meanwhile, it is well-known that the Kretschmann geometry can efficiently excite the




FIG. 1: (Color online) Optical forces at different height from the prism as functions of the wavelength. (Blue) dotted line: a microcylinder illuminated by a plane wave in the absence of the prism. (Black) solid line: the horizontal force F_x acting on a microcylinder with the prism. (Red) dashed line: the vertical force F_y with the prism. (a) $d = 1.1a$, (b) $d = 1.2a$, (c) $d = 1.3a$, Inset of (c): an enlarged section of (c). (d) $d = 1.1a$ and (e) $d = 1.2a$ for shorter wavelength. The Q_{iso} and resonant wavelength for the typical modes in (d) and (e) are, respectively, 5399 and $0.31110a$ for TE_{26}^1 , 7632 and $0.30047a$ for TE_{27}^1 , 10817 and $0.29055a$ for TE_{28}^1 , 15368 and $0.28128a$ for TE_{29}^1 , 96 and $0.31044a$ for TE_{22}^2 , 115 and $0.29928a$ for TE_{23}^2 , 139 and $0.28892a$ for TE_{24}^2 , and 170 and $0.27926a$ for TE_{25}^2 . (f) Illustration of the Kretschmann geometry, where the evanescent wave is generated by total internal reflection on top of the prism.

*Electronic mail: jjxiao@ust.hk

†Electronic mail: jack@ust.hk

WGM of a high- Q microsphere. Kawata and Sugiura³ demonstrated versatile optical manipulation by evanescent waves subsequently.²¹ Although the Kretschmann geometry is well-known to be suitable for both optical manipulation and the excitation of WGM, the resonant forces associated with evanescent wave excited WGM have never been considered experimentally. Here, we hope to provide a more complete treatment for the WGM induced forces, and hope our work can stimulate experimental works in this area.

In Figs. 1(a)–1(c), we show the optical forces acting on a MC for $\lambda = 0.32a - 0.4a$ (size parameter $k_p a \approx 24 - 30$). The (blue) dotted line represents both the horizontal and vertical forces (equal by symmetry) acting on a bare MC without the prism.²² The (black) solid line represents the horizontal force F_x and the (red) dashed line represents the vertical force F_y , in the presence of the prism. For the case without the prism, the plane incident wave hits directly on the MC, and for the case with the prism, the plane wave is partially reflected by the prism surfaces and partially converted into an evanescent wave on the prism's top surface. In the case of a bare MC, it can be seen from Fig. 1(a) that the force at resonance is comparable to the force at off resonance; thus size-selectivity is weak. On the other hand, the case with an incident evanescent wave (generated by the prism) is very size-selective, as we shall see below.

Let us first consider the horizontal force F_x . As shown in Fig. 1(c) where the separation $d = 1.3a$, it is quite clear that the force on first radial order resonance is significantly stronger than the force at off resonance, and force on the second order resonances can hardly be seen at all. This results in a high peak-to-baseline ratio, defined as the peak resonance force divided by the maximum of off resonance or lower- Q resonance force. This would imply the possibility of size-selective manipulation: upon illumination, only those particles whose sizes happen to be in resonance with the incident wave will experience a sizable optical force, whereas the off resonance particle will be left untouched. In the limit of large separation, the peak-to-baseline ratio can be several orders of magnitude higher than what is predicted here, as found out in Ref. 2. Indeed, the results presented in Fig. 1(c) already begin to resemble that of Ref. 2.

Although it is highly desirable to have a high peak-to-baseline ratio, it is also important to achieve large force. This case is illustrated in Figs. 1(a) and 1(b). As the separation decreases from $d = 1.3a$ in Fig. 1(c) to $d = 1.2a$ in Fig. 1(b), the force on first order resonance increases by a factor of $\sim 2-3$ as the intensity becomes higher in the near field, while its linewidth is broadened slightly. At $d = 1.2a$, the second order resonance can now be observed, due to the higher intensity. Although the peak-to-baseline ratio for $d = 1.2a$ is smaller than that of $d = 1.3a$, the size selectivity is still very strong. At an even closer separation [$d = 1.1a$ in Fig. 1(a)], although the intensity of the evanescent wave increases, the strength of the first order resonance force remains at the same level as when $d = 1.2a$. This is a consequence of the Lorentzian lineshape,² which implies that the resonance force is inversely proportional to the linewidth, or directly proportional to its Q . As d decreases, the stronger modal coupling between the prism and the MC broadens the linewidth (or decrease in Q), and causes the resonance force to decrease, compensating the increase in the local in-




FIG. 2: (Color online) Optical forces as a function of cylinder-prism separation d at various situations. (a) at the TE_{22}^1 resonance, (b) at an off resonance frequency between TE_{22}^1 and TE_{18}^2 , (c) at the TE_{18}^2 resonance.

tensity I_{loc} . The background force and the forces associated with the second order resonance have increased considerably, making the peak-to-baseline ratio small. We note that as the MC approaches the prism, the resonance frequencies do not shift and it is the Q 's that drops. This decrease in Q will significantly reduce the size-sensitivity of the resonance force, which scales like $\sim 1/Q$.

A more explicit demonstration on the selectivity due to high- Q is shown in Fig. 1(d) and 1(e) which show the optical forces with the presence of the prism in higher size parameter regime $\lambda = 0.28a - 0.32a$ ($k_p a \approx 30 - 34$). It is seen in Fig. 1(d) that with $d = 1.1a$ several radially second order WGMs can be well excited, resulting in comparable optical forces of the radially first order WGMs as indicated by the downward arrows. We find that the Q value of the TE_{23}^2 mode exceeds 10^2 (see the details in the caption of Fig. 1). The MC with such a value of Q can already collect sufficient incident field energy that yields a non-negligible amount of light scattering. Figure 1(e) shows that for increased separation to $d = 1.2a$, these second order modes remain not well excited again, simply because of the decreased local field. At this separation, however, the radially first order WGMs with Q in the order of 10^3 and 10^4 can be well excited and are manifested as the several sharp peaks [as labeled in Fig. 1(e)].

Figure 2 clarifies the separation dependence of the optical force on and off resonance. Here we have chosen three different frequencies, one for TE_{22}^1 (with quality factor of isolated particle $Q_{\text{iso}} = 1395$)¹⁰ shown in Fig. 2(a), one for TE_{18}^2 ($Q_{\text{iso}} = 49$)¹⁰ shown in Fig. 2(c), and the last one for an off resonance frequency shown in Fig. 2(b). We remark that the strength of the resonance force depends on both the local intensity I_{loc} and the mode's Q , i.e. $F \propto Q I_{\text{loc}}$. The observed Q in the force spectrum is in fact related to the original quality factor for the isolated MC, Q_{iso} , and the coupling Q -factor, Q_{coup} , through the equation $1/Q = 1/Q_{\text{iso}} + 1/Q_{\text{coup}}$. In Fig. 2(b) and 2(c), the optical force decreases monotonically as d increases. The off resonance case of Fig. 2(b) is straightforward to understand: as d increases, the local intensity decreases, so the force decreases as well. For the second order resonance TE_{18}^2 case plotted in Fig. 2(c), the system's Q is completely determined by its




FIG. 3: (Color online) Near fields for high- Q (left column) and low- Q (right column) resonances. (a) and (c) are, respectively, for TE_{22}^1 and TE_{18}^2 in the absence of the prism. (b) and (d) are, respectively, for the same resonances as (a) and (c), but with the prism at a separation $d = 1.2a$ (lower panels).

small $Q_{\text{iso}} = 49$ (because $Q_{\text{iso}} \ll Q_{\text{coup}}$), consequently as d increases, the total Q is essentially unchanged while I_{loc} decreases, causing the optical force to drop. For the high- Q resonance of TE_{22}^1 shown in Fig. 2(a), it can be anticipated that the optical forces will be weak when the MC is either too close to (due to a small Q) or far away from (due to small I_{loc}) the prism, and there is a peaked force in between. In Fig. 2(a), this peak occurs at $d \approx 1.16a$.

To give an overall perspective of the effect of Q value and the effect of the prism, we compare the near fields in Fig. 3 for four different situations corresponding to Figs. 2(a) and 2(c). These figures clearly demonstrate the resonant tunneling mechanism which accounts for the optical forces. We can draw a conclusion that a high Q value is the prerequisite for the resonant tunneling. Therefore in the scheme with the prism for optical sorting operated at a specific frequency, high Q particles can be easily distinguished and selected.

We finally consider the vertical force F_y . As Rayleigh dielectric objects tend to move to region of high intensity in order to minimize the electromagnetic free energy, one may believe that the vertical force is attractive relative to the prism. But as shown in Fig. 1(c) where $d = 1.3a$, the vertical force versus wavelength curve actually consists of a background repulsion which lifts the entire curve upward plus a peak and dip pair at each resonance position. As the MC approaches the prism [Figs. 1(a) and 1(b)], the line-

shape of F_y evolves and deviates from that of the original lineshape in Fig. 1(c). We see that the resonance force has become stronger, and F_y can now be either attractive (at the dip) or repulsive (at the peak). This resonance lineshape (the peak and dip pair) can be understood as follows: when the frequency changes across the resonance frequency, the WGM switches from in phase to out of phase with the incident field, and this results in a change of the sign in the force. We note that in addition to typical resonance peak-and-dip, there is still a bias of the force to the repulsive side. Heuristically, this repulsive bias can be traced to the fact that the incident photon momentum is upward, so it should not be surprising to see the photons pushing the MC up. We note that in the 1992 experiment of Kawata and Sugiura,³ a repulsive force is also observed. This subtle effect warrants further studies.

In summary, we have numerically investigated the WGM enhanced optical force as excited by an evanescent wave. The high peak-to-baseline ratio associated with evanescent wave excitation makes it highly size-selective and may thus be useful for particle sorting. The notion of Ref. 2 is confirmed even in the presence of a substrate, and we have gone beyond Ref. 2 and show that the idea even works in the substrate's near field. In general, a high Q mode guarantees a stronger optical force, as it favors the resonant tunneling of the evanescent light. As a consequence of the competition between the increased coupling loss at small separation and the decreased intensity at large separation, there exists an optimal particle-prism separation in which the resonance force is maximized. Accordingly, in an optical sorting experiment, it would be highly desirable to have some other means to keep the particles at an appropriate height, for if the MC is too close to the prism surface, the linewidth broadening would reduce the sorting accuracy as well as the peak-to-baseline ratio, but if the MC is too far away, the force is too weak. Finally, we remark that the induced optical force can further be enhanced by coating the substrate with a metallic coating that supports surface plasmon,²³ or by coating the prism with a dielectric cavity layer.²⁴

This work was supported by the Hong Kong RGC Grant No. 600308. ZFL was also supported by NSFC (10774028), PCSIRT and MOE of China (B06011). Computational resources were supported by the Shun Hing Education and Charity Fund. JJX acknowledges Jeffrey C. W. Lee's technical help on the MPI.

- ¹ *Optical Microcavities*, edited by K. J. Vahala (World Scientific, Singapore, 2004), Chap 5.
- ² J. Ng and C. T. Chan, *Appl. Phys. Lett.* **92**, 251109 (2008).
- ³ S. Kawata and T. Sugiura, *Opt. Lett.* **17**, 772 (1992).
- ⁴ S. Chang, J. T. Kim, J. H. Jo, and S. S. Lee, *Opt. Commun.* **139**, 252 (1997).
- ⁵ M. Lester and M. Nieto-Vesperinas, *Opt. Lett.* **24**, 936 (1999).
- ⁶ M. Gu, J. -B. Haumonte, Y. Micheau, J. W. M. Chon, and X. Gan, *Appl. Phys. Lett.* **84**, 4236 (2004).
- ⁷ H. Y. Jaising and O. G. Helleso, *Opt. Commun.* **246**, 373 (2004).
- ⁸ Other schemes of optical sorting were also proposed; see e.g. P. T. Korda, M. B. Taylor, and D. G. Grier, *Phys. Rev. Lett.*

- 89**, 128301 (2002), M. P. Macdonald, G. C. Spalding, and K. Dholakia, *Nature (London)* **426**, 421 (2003), and M. Righini, A. S. Zelenina, C. Girard, and . Quidant, *Nat. Phys.* **3**, 477 (2007).
- ⁹ Y. Hara, T. Mukaiyama, K. Takeda, and M. Kuwata-Gonokami, *Opt. Lett.* **28**, 2437 (2003).
- ¹⁰ J. J. Xiao and C. T. Chan, *J. Opt. Soc. Am. B* **25**, 1553 (2008).
- ¹¹ J. Ng, Z. F. Lin, C. T. Chan, and P. Sheng, *Phys. Rev. B* **72**, 085130 (2005).
- ¹² P. Chylek, J. T. Kiehl, and M. K. W. Ko, *Phys. Rev. A* **18**, 2229 (1978).
- ¹³ E. Almaas, and I. Brevik, *J. Opt. Soc. Am. B* **12**, 2429 (1995).
- ¹⁴ J. R. Arias-Gonzalez and M. Nieto-Vesperinas, *Opt. Lett.* **25**,

- 782 (2000).
- ¹⁵ T. Carmon, H. Rokhsari, L. Yang, T. J. Kippenberg, and K. J. Vahala, Phys. Rev. Lett. **94**, 223902 (2005).
 - ¹⁶ J. Ng, C. T. Chan, P. Sheng, and Z. F. Lin, Opt. Lett. **30**, 1956 (2005).
 - ¹⁷ M. Eichenfield, C. P. Michael, R. Perahia, and O. Painter, Nat. Photonics **1**, 416 (2007)
 - ¹⁸ A. B. Stilgoe, T. A. Nieminen, G. Knoner, N. R. Heckenberg, and H. Rubinsztein-Dunlop, Opt. Express **16**, 15039 (2008).
 - ¹⁹ A. Ashkin, Science **210**, 1081 (1980).
 - ²⁰ A. Fontes, A. A. R. Neves, W. L. Moreira, A. A. de Thomaz, L. C. Barbosa, C. L. Cesar, and A. M. de Paula, Appl. Phys Lett. **87**, 221109 (2005).
 - ²¹ K. Dholakia and P. Reece, *Near-Field Optical Micromanipulation*, in *Structured Light and Its Applications*, edited by D. L. Andrews (Academic Press, Amsterdam, 2008).
 - ²² Our calculation has been checked against the semi-analytical result of Grzegorzczuk and Kong, and excellent agreement was achieved. See e.g. T. M. Grzegorzczuk and J. Kong, J. Opt. Soc. Am. B **24**, 644 (2007) and *ibid*, Opt. Express **13**, 8010 (2007).
 - ²³ V. Garces-Chavez, R. Quidant, G. Badenes, and D. Petrov, Phys. Rev. Lett. **96**, 238101 (2006).
 - ²⁴ P. J. Reece, V. Garces-Chavez, and K. Dholakia, App. Phys. Lett. **88**, 221116 (2006).